

MAVERICE[®]

insecticide



SPECIMEN

ROACH CONTROL SYSTEM™

FOR USE IN COMMERCIAL, INDUSTRIAL AND RESIDENTIAL AREAS

Active Ingredient:

Hydramethylnon [Tetrahydro-5,5-dimethyl-2 (1H)-pyrimidinone (3-[4-(trifluoromethyl)phenyl]-1-(2-[4-(trifluoromethyl)phenyl]ethenyl)-2-propenylidene)hydrazone]	1.65%
Inert Ingredients	98.35%
Total	100.00%

PRECAUCIÓN

AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel For Additional Precautions



© 1987 American Cyanamid Company
Household Products
Wayne, NJ 07470

Total Net Weight: 3.8 oz. (108 g)

EPA Reg. No. 1730-67
EPA Est. No. 1730 MS-01
Product Code No. 24563-02

T 141 F

CODE 72250

DRIONE[®] Insecticide

EPA Reg. No. 4816-353 EPA Est. No. 279-111-1

A Delicant Dust
PROVIDES QUICK CONTROL FOR INDUSTRIAL COMMERCIAL USE ONLY. (TUB LARGER THAN 4 OZ. CONTAINERS ONLY)
KILLS UP TO 6 MONTHS WHEN LEFT UNDISTURBED.
NON-STAINING AND RELATIVELY ODRPLESS.
FOR USE IN RESIDENCES, INSTITUTIONS, FOOD PLANTS, GRANARIES AND ON CATS AND DOGS.
CONTROLS ANTS, COCKROACHES, SILVERFISH, BEDBUGS, FLEAS, LICE, BEES, AND AS STIPULATED IN THE USE DIRECTIONS ON NUMEROUS STORED PRODUCT PESTS. TO PROTECT AGAINST ENTRANCE OF DRYWOOD TERMITES.

ACTIVE INGREDIENTS:

Pyrethrins.....	1.0%
Piperonyl Butoxide, Technical.....	10.0%
Amorphous Silica Gel.....	40.0%
INERT INGREDIENTS:	49.0%
	100.0%

*Equivalent to 8.0% (butylcarbityl) (6-propylpiperonyl) ether and 2.0% related compounds.
**Contains Petroleum Distillate.

KEEP OUT OF REACH OF CHILDREN.
CAUTION
See back panel for additional precautionary statements.

DRIONE - Registered Trademark of Fairfield American Corporation.

NET CONTENTS:
BATCH

Mixed in U.S.A.

FAIRFIELD AMERICAN CORPORATION
201 ROUTE 17 NORTH - NORTHWOOD, NJ 07070

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

APPLICATIONS WITHIN FOOD AREAS:

Food processing plants and warehouses, bakeries, beverage plants, restaurants, supermarkets, meat and poultry processing plants, hotels, motels, hospitals, and schools:
Limit to Duster and Crevice Treatment Only:
Using a bulb-duster apply a small amount of material directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids, motor housing, junction boxes or switch boxes, conduits or hollow equipment legs where cockroaches, ants, silverfish, spiders, boxelder bugs, and crickets hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT, ARE NOT PERMITTED.

NON-FOOD AREAS:

Homes, institutions, hospitals, office buildings, hotels, motels, theaters, schools, warehouses, truck trailers and railroad cars: To control crawling insects such as ants, cockroaches, silverfish, firebrats, spiders, boxelder bugs, ticks, lice, and fleas in animal quarters, distribute Dione at the rate of 2 ounces per 100 square feet of surface area. Apply a continuous visible film underneath and behind stairs, refrigerators and appliances. Treat on, around and behind moldings, shelving, baseboards, pipe openings, cracks, crevices, and other areas where insects hide. For severe infestations, apply Dione behind cabinets and walls, in crawl spaces and attics at the rate of 16 ounces per 1000 square feet of surface area. Repeat treatment as necessary.

STORED PRODUCT PESTS:

To aid in the control of cadellies, cheese mites, cigarette beetles, confused flour beetles, dark meal worms, drugstore beetles, grain mites, grainy weevils, red flour beetles, rice weevils, sawtoothed grain beetles and yellow meal worms in mills, elevators and granaries. Clean up waste materials, dust, dirt, and any other debris. Thoroughly treat floors, walls and other surfaces in bins, storage and handling areas. Apply into cracks and crevices and other difficult to reach areas where insects may hide.

BEES AND WASPS:

It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Using hand or power duster or other suitable means, with extension tubes if necessary, thoroughly dust nest and entrance and surrounding areas where insects alight. Nests in wall voids can be located by listening with your ear against the wall. Drill a hole in the wall.

emergence of newly-hatched insects. If removal is not feasible, retreat the nest, if necessary. Do not use this product to treat overhead nests.

SEWER TREATMENTS:

American and Oriental cockroach infestations in sanitary sewers may be treated with approximately 6 ounces of Dione per manhole. When treating sewers, one manhole per 200 feet of line should be treated. Treatments are made by removing the manhole cover and replacing with a disc of plywood bored to receive the usual flexible hose from portable to truck-mounted ventilating equipment. 6 ounces of Dione is introduced into the ventilating air to be dispersed quickly and thoroughly through the system.

BEDBUGS:

Take bed apart. Dust into joints and channels. If hollow, such as square or round tubing, see that the interior of framework is well dusted. Mattresses should be dusted, especially tufts, folds, and edges. Picture frame moldings and all crack and crevices in the room should be treated.

DRYWOOD TERMITES:

To protect against entrance, using an electric or hand operated rotary duster, distribute Dione through an entry, e.g. crawl hole into attic and crawl space of building, at a rate of 1 pound per 1000 square feet of floor area. All vents and blowers on the windward side should be closed. All wood surfaces should be covered. In case of localized infestations probe to locate galleries and drill holes to inject Dione into the galleries. If a reasonably heavy infestation has already developed, fumigation should be employed.

FLEAS AND TICKS ON DOGS AND CATS:

Liberaly apply powder to the animal. Rubbing thoroughly to the skin. Begin with the neck and include the underbody, legs, feet and the base of the tail. Repeat as necessary. Surfaces of kennels should be dusted at the rate of one ounce per 50 square feet. Heavily infested lawns, basements, and crawl spaces should be dusted at the rate of 16 ounces per 1000 square feet. Repeat as necessary.

Do NOT treat kittens or puppies less than four weeks old.

SOW BUGS, PILLBUGS, GROUND BEETLES, EARWIGS, MILLPEDES, CENTIPEDES, AND SCORPIONS:

Apply around doors and windows and other places where these pests may enter premises. Treat baseboards, storage areas and other locations where the pests are found.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Keep away from moisture. Sweep up and dispose of with chemical waste. PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that